

Losartan

Angiotensin II is a substance that causes the blood vessels to become narrower and this causes an increase in blood pressure. Losartan lowers blood pressure by blocking the effects of a substance called Angiotensin II at the blood vessel walls. This medication can also be used for patients who have protein in their urine and other heart conditions.

What is this medication for?

Angiotensin II is a substance that causes the blood vessels to become narrower and this causes an increase in blood pressure. Losartan lowers blood pressure by blocking the effects of a substance called Angiotensin II at the blood vessel walls.

This medication can also be used for patients who have protein in their urine and other heart conditions.

How should I take/use this medication?

- Do not stop taking your medicine without checking with your healthcare provider
- This medicine may be taken with or without food

What should I do if I forget to take/use this medication?

- If you miss a dose, take the missed dose as soon as you remember. If it is almost time for your next dose, take only the usual dose. Do not double the dose or use extra medicine to make up for the missed dose.

What precautions should I take?

Inform your doctor if you are pregnant or planning to get pregnant. Once you find out you are pregnant, stop the medication as soon as possible. Taking this medication can cause injury and death to your developing fetus.

Your doctor may need to monitor your potassium levels and kidney function when you are on this medication.

What are some side-effects that I need to seek medical advice immediately?

The symptoms of a drug allergy include one or more of the following:

- Swollen face/eyes/lips/tongue
- Difficulty in breathing
- Itchy skin rashes over your whole body

If you experience any of these symptoms, you should stop your medicine immediately and see your doctor.

Very rarely, this medicine can injure or damage the liver. A liver injury may recover on its own. However in some cases, it can cause serious liver damage. If you are taking this medicine and see a doctor immediately if you notice dark coloured urine or light coloured stools, or if you experience nausea, vomiting, loss of appetite, stomach pain or yellowing of your eyes or skin. This side effect has a very low chance of occurring. Most patients take this medicine with no problems. Tell your doctor or pharmacist if you have any concerns regarding your medicines.

What are some other side-effects of this medication?

The side effects include dizziness and lightheadedness on getting up.

How should I store this medication?



- Store your medication in a cool, dry place away from heat, moisture and direct sunlight.
- Keep the medicine out of reach of children.
- Throw away all expired medicines.

How should I dispose of this medication safely?

Pack this medicine into a black trash bag and seal it tightly before throwing into the rubbish chute or bin.

Disclaimers

If you take more than the recommended dose, please seek medical advice immediately. The information provided on this page does not replace inform your healthcare professional. Please consult your healthcare professional for more information.

Community Partners



Cluster Partners



Brought to you by



This article was last reviewed on Tuesday, April 30, 2019

