

Umeclidinium

Medication Information Leaflet

What is this medication for?

Umeclidinium inhaler is used for the long term treatment of people who have chronic obstructive pulmonary disease (COPD). COPD is a disease affecting the lungs which makes it harder for the person to breathe as the disease gets worse.

Umeclidinium works by opening up the airways and make it easier for air to move in and out of the lungs.

How should I take/use this medication?

- Use your Umeclidinium inhaler every day to control your condition.
- Do not stop using your inhaler without checking with your healthcare professionals.
- Please refer to the Umeclidinium inhaler's patient information leaflet to learn how to use the inhaler.

What should I do if I forget to take/use this medication?

If you forget to take a dose, take it as soon as you remember. Then take your next dose at the usual time. Do not take two doses to make up for the missed dose.

What precautions should I take?

Inform your healthcare professional if:

- You are allergic to this medication or any of the other ingredients of this medication
- You are pregnant, planning to become pregnant, or breastfeeding
- You are taking any other medications, including supplements, traditional medications and herbal remedies
- You have the following medical conditions:
 - Narrow-angle glaucoma (a type of eye condition where the pressure in the eye is increased)
 - Prostate problems or difficulty passing urine
 - Heart conditions such as irregular heart beats
 - Serious liver problems
 - Lactose intolerance (a condition where your body is unable to break down a type of sugar found naturally in milk called lactose)

What are some side-effects that I need to seek medical advice immediately?

In very rare cases, some patients might experience the following:

- Difficulty in passing urine
- Eye pain, blurred vision, red eyes
- Sudden difficulties in breathing such as shortness of breath, wheezing (a high-pitched whistling sound made while breathing) or chest tightness

The symptoms of a drug allergy include one or more of the following:

- Swollen face/eyes/lips/tongue
- Difficulty in breathing
- Itchy skin rashes over your whole body

If you experience any of these symptoms, you should stop your medication and see your healthcare professional immediately.

What are some other side-effects of this medication?

Common side effects include:

- Symptoms of upper respiratory tract infections such as cough, runny nose, sore throat and pain at the cheek and forehead areas

- Painful and frequent urination, which may be signs of a urinary tract infection
- Fast heart beat
- Dry mouth
- Nausea
- Constipation
- Headache

These symptoms may happen but should get better over time. Check with your doctor if any of these symptoms are serious or do not go away.

How should I store this medication?

Store in a cool and dry place, away from direct sunlight. Keep this medication away from children. Keep the cap on when not in use.

When should I throw away the medication?

Check the dose counter and change the inhaler once the dose counter indicates zero doses or six weeks after you opened it.

How should I throw away this medication safely?

Pack this medication into a black trash bag and seal it tightly before throwing into the rubbish chute or bin.

Disclaimers

If you take more than the recommended dose, please seek medical advice immediately. The information provided on this page does not replace information from your healthcare professional. Please consult your healthcare professional for more information.

This article is jointly developed by members of the National Medication Information workgroup. The workgroup consists of cluster partners (National Healthcare Group, National University Health System and SingHealth), community pharmacies (Guardian, Unity and Watsons) and Pharmaceutical Society of Singapore.

Last updated on July 2019

This article was last reviewed on Monday, September 2, 2019