Medication information leaflet



[Dequalinium Lozenges]

What is this medication for?

Dequalinium lozenges have an antiseptic effect and is used for the symptomatic relief of sore throat.

How should I take/use this medication?

- Suck Dequalinium lozenges according to the instructions on the medicine label
- You may take this medication with or without food

What should I do if I forget to take this medication?

If you forget to take a dose, take it as soon as you remember. Then take your next dose at the usual time. Do not take two doses to make up for the missed dose.

What precautions should I take?

Inform your healthcare professional if:

You are allergic to this medication or any of the other ingredients of this medication

What are some common side-effects of this medication?

- Dequalinium lozenges are generally free of side effects when taken at recommended doses
- Sometimes it may cause soreness of the tongue, mouth discomfort or a burning sensation in the mouth
 - Stop sucking Dequalinium lozenges if that happens

What are some rare but serious side-effects that I need to seek medical advice immediately? The symptoms of a drug allergy include one or more of the following:

- Swollen face/eyes/lips/tongue
- Difficulty in breathing
- Itchy skin rashes over your whole body

If you experience any of these symptoms, you should stop your medication and see your healthcare professional immediately.

How should I store this medication?

Store in a cool and dry place, away from direct sunlight. Keep this medication away from children.

Last updated on October 2021