
PATIENT INFORMATION LEAFLET

ANTIVIRALS







V

Alexandra Health EASTERN HEALTH ALLIANCE

Jurong Health



SINGHEALTH / PIL / E66R0

SingHealth

ABOUT your medicine

Acyclovir and Valaciclovir are antiviral drugs. They are used to treat infections caused by herpes viruses including genital herpes, cold sores, shingles and chicken pox. They slow the growth and spread of the herpes virus so that the body can fight off the infection.

Valaciclovir is a prodrug and it will be converted into the active drug acyclovir by the liver.

Acyclovir and Valaciclovir may also be used to treat other conditions.

HOW should I take the medicine?

Take Acyclovir or Valaciclovir exactly as directed by your doctor. Take each dose with a full glass of water. Acyclovir and Valaciclovir can be taken with or without food. Taking Acyclovir or Valaciclovir with food may decrease stomach upset.

Suspensions should be shaken well before measuring a dose. To ensure that you get a correct dose, measure the suspension with a special dose-measuring spoon or cup, instead of a regular table spoon.

What should I do if I FORGET a dose?

If you miss a dose or forget to take your medicine, take it as soon as you can. However, if it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not double your dose or take extra medicine to make up for a missed dose. If you have missed several doses, please call your physician to check what you should do.

• Unusual bleeding or bruising

Other less serious side-effects may more likely occur. Continue to take Acyclovir or Valaciclovir and talk to your doctor if you experience:

What are the possible SIDE-EFFECTS of the medicine?

Stop taking Acyclovir or Valaciclovir and seek emergency medical attention or contact

your doctor immediately if you experience any of the following serious side-effects:

Nausea

Hives

•

- Vomiting
- Diarrhoea
- Decreased appetite

· Difficulty breathing

Closing of your throat

Little or no urine production

Swelling of your lips, tongue or face

- Abdominal pain
- Headache or light-headedness
- Joint pain

Side-effects other than those listed above may also occur. Talk to your doctor about any side-effect that seems unusual or that is especially bothersome.

What PRECAUTIONS should I take?

Inform your doctor or pharmacist if you are already taking some other medicines, herbal preparations, nutritional products or vitamin supplements.

Inform your doctor or pharmacist if you have kidney disease.

Inform your doctor or pharmacist if you are allergic to any other medicines.

Inform your doctor if you are breast-feeding, pregnant, or likely to become pregnant during the course of the treatment.

How should I STORE the medicine?

Store the medicine in a dry place at room temperature, away from heat and direct light. Do not refrigerate your medicines unless instructed to do so.